



## VACCINES REQUIRED FOR SCHOOL ATTENDANCE, GRADES K-12

### August 1, 2020 – July 31, 2021

VACCINE	Kindergarten - 6 <sup>th</sup> Grade	7 <sup>th</sup> – 9 <sup>th</sup> Grade	10 <sup>th</sup> - 12 <sup>th</sup> Grade
<b>Hepatitis B</b>	<b>3 doses</b> Dose 3 must be given on or after 24 weeks of age		
<b>DTaP</b> (Diphtheria, Tetanus, and Pertussis)	<b>5 doses</b> (4 doses only <b>IF</b> 4 <sup>th</sup> dose given on or after 4 <sup>th</sup> birthday <b>AND</b> a minimum interval of 6 months from the previous dose)		
<b>Tdap</b> (Diphtheria, Tetanus, and Pertussis)	<b>Not Required</b>	<b>1 dose Tdap</b> <i>(see page 2 for more details)</i>	
<b>IPV</b> (Polio, for OPV see page 2)	<b>4 doses</b> (3 doses only <b>IF</b> 3 <sup>rd</sup> dose given on or after 4 <sup>th</sup> birthday) The final dose given on or after August 7, 2009, must be given on or after 4 years of age <b>AND</b> a minimum interval of 6 months from the previous dose.		<b>4 doses</b> (3 doses only <b>IF</b> 3 <sup>rd</sup> dose given on or after 4 <sup>th</sup> birthday)
<b>MMR</b> (Measles, Mumps, and Rubella)	<b>2 doses</b>		
<b>Varicella</b> (Chickenpox)	<b>2 doses</b> <b>OR</b> Healthcare provider verified disease history		

- Look at the Minimum Age and Interval Table on page 2 for recommended minimum age and spacing information.
- Review the Individual Vaccine Requirements Summary for more detailed information: <https://www.doh.wa.gov/SCCI>.

## Minimum Age & Interval for Valid Vaccine Doses

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
Hepatitis B <b>HepB</b>	Dose 1	Birth	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>2 doses acceptable if both doses are documented as adult dose of Recombivax HB® given between ages 11 and 15. The doses must be separated by at least 4 months.</li> </ul>
	Dose 2	4 weeks	8 weeks between Dose 2 & 3	
	Dose 3	24 weeks	16 weeks between Dose 1 & 3	
Diphtheria, Tetanus, and Pertussis <b>DTaP</b>	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>6 month interval is recommended between Dose 3 and Dose 4, but a minimum interval of 4 months is acceptable.</li> <li>DTaP: can be given to children through age 6. If catch-up doses are needed at age 7 and older Tdap is used followed by additional doses of Td if needed. A Tdap given at age 7 through 9 years of age does not count for the 7<sup>th</sup> grade Tdap requirement.</li> <li>See the Individual Vaccine Requirements Summary for more details about the catch-up schedules: <a href="https://www.doh.wa.gov/SCCI">https://www.doh.wa.gov/SCCI</a></li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	12 months	6 months between Dose 4 & 5	
	Dose 5	4 years	–	
Tetanus, Diphtheria, and Pertussis <b>Tdap</b>	Dose 1	11 years See notes for exceptions	–	<ul style="list-style-type: none"> <li>Tdap booster dose recommended at age 11 is required for all students in grades 7-12.</li> <li>For students in 7<sup>th</sup> grade: Tdap dose acceptable if given on or after 10 years of age.</li> <li>For students in 8<sup>th</sup>-12<sup>th</sup> grades: Tdap dose acceptable if given on or after 7 years of age.</li> <li>Tdap booster dose can be given regardless of the interval between DTaP/DT/Tdap/Td.</li> </ul>
Polio <b>IPV or OPV</b>	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>OPV given on or after 04/01/16 cannot be accepted as a valid dose in the series.</li> <li>Not required for students 18 years and older.</li> <li>Please see Individual Vaccine Requirements Summary for more details: <a href="https://www.doh.wa.gov/SCCI">https://www.doh.wa.gov/SCCI</a></li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	4 years	–	
Measles, Mumps, and Rubella <b>MMR</b>	Dose 1	12 months	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>MMRV (MMR + varicella) may be used instead of separate MMR and varicella vaccines.</li> <li>Must get the same day as other live vaccine (ex. varicella, Flumist) <b>OR</b> at least 28 days apart.</li> <li>4-day grace <b>DOES</b> apply between doses of the same live vaccine such as MMR/MMR or MMRV/MMRV. The 4 day grace period <b>DOES NOT</b> apply between dose 1 and dose 2 of different live vaccines, such as between MMR and Varicella or between MMR and live flu vaccine.</li> </ul>
	Dose 2	13 months	–	
Varicella (chickenpox) <b>VAR</b>	Dose 1	12 months	3 months between Dose 1 & 2 (12 months through 12 years) 4 weeks between Dose 1 & 2 (13 years and older)	<ul style="list-style-type: none"> <li>Age 12 months through 12 years: 3 months between varicella doses recommended, but minimum interval of 28 days acceptable on retrospective record review.</li> <li>MMRV (MMR + varicella) may be used instead of separate MMR and varicella vaccines.</li> <li>Must get the same day as other live vaccine (ex. MMR, Flumist) <b>OR</b> at least 28 days apart.</li> <li>4-day grace period <b>DOES</b> apply between doses of the same live vaccine such as varicella/varicella or MMRV/MMRV; The 4 day grace period <b>DOES NOT</b> apply between doses of different live vaccines, such as between MMR and varicella or between varicella and live flu vaccine.</li> </ul>
	Dose 2	15 months	–	